

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.

TITLE: OXIDATION OF HETEROCYCLIC AMINES, SULFILIMINES,
AND PHOSPHINE IMINES WITH DIMETHYLDIOXIRANE

AUTHOR(S): Michael D. Coburn

LA-UR--87-1739

DE87 010129

SUBMITTED TO: Sixth Annual Working Group Meeting on
Synthesis of High Energy Density Materials
Kiamesha Lake, NY
May 19-22, 1987

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U S Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U S Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U S Department of Energy

 Los Alamos National Laboratory
Los Alamos, New Mexico 87545

OXIDATION OF HETEROCYCLIC AMINES, SULFILIMINES, AND
PHOSPHINE IMINES WITH DIMETHYLDIOXIRANE

Michael D. Coburn
University of California, Los Alamos National Laboratory
P.O. Box 1663, Los Alamos, NM 87545

ABSTRACT

Our previous work¹ has shown that oxidation of 3-amino-4-(4-chlorophenyl)furazan (1) with peroxytrifluoroacetic acid (ptfa) gave azoxy (4-chlorophenyl)furazan (6) as the major product along with a small amount of 3-(4-chlorophenyl)-4-nitrofurazan (5). The dimethylsulfilimine 2 derived from 1 gave near quantitative yields of 5 when subjected to oxidation with either ptfa or m-chloroperoxybenzoic acid (mcpba). In contrast, both the trioctylphosphine imine 3 and the triphenylphosphine imine 4 derived from 1 were oxidized by mcpba to give 6 as the exclusive product (Fig. 1).

It was recently reported that dimethyldioxirane (DMD), which is conveniently prepared by oxidation of acetone with persulfate², will convert certain aromatic amines to nitro compounds in high yield. In contrast to our results with peracids, DMD reacted with 1 to yield 5 as the exclusive product, but with 2 to give the sulfoximine (7) as the major product with only a small amount of 5. However, treatment of 3 with DMD gave 5 as the exclusive product (Fig. 1).

¹M. D. Coburn, J. Heterocyclic Chem. 23, 421 (1986).

²R. W. Murray and R. Jeyaraman, J. Org. Chem. 50, 2847 (1985).

3-Amino-4-nitrofurazan (8) failed to react with DMD and its sulfilimine (9) and phosphine imine (10) derivatives gave the oxidized species 11 and 12 (Fig. 2). We were surprised to find that 4,4'-diamino-3,3'-bifurazan (13) would not react with DMD, since it is readily oxidized to 4,4'-dinitro-3,3'-bifurazan with ptfa.³


2-Aminopyridine (14) was converted to its N-oxide (15) with DMD and its sulfilimine (16) and phosphine imine (17) derivatives were oxidized smoothly to 2-nitropyridine (18) with DMD (Fig. 3).

Both 2-aminopyrimidine (19) and 2-amino-5-nitropyrimidine (20) gave the N-oxides 21 and 22 when treated with DMD. The sulfilimine derivative of 20 (23) was converted with DMD to the sulfoximine (24). Although not isolated, 2-nitropyrimidine may have formed when the phosphine imine of 19 (25) was treated with DMD (Fig. 4).

Treatment of 2-aminopyrazine (26) with DMD produced the 1,4-dioxide (27). The phosphine imine derivative (28) was converted to a mixture of oxygenated products, possibly 29 and 30; however, the sulfilimine (31) was smoothly oxidized to 2-nitropyrzine (32) with DMD (Fig. 5).

Oxidation of 3,6-diamino-1,2,4,5-tetrazine (33) with DMD gave the 1,4-dioxide (34), which is predicted to have a crystal density of 1.95 g/cm³ by J. R. Stine. Treatment of the disulfilimine (35) derived from 33 with mcpba gave what appears to be 3-(S,S-dimethylsulfilimino)-6-nitroso-1,2,4,5-tetrazine (36), which was subjected to oxidation by DMD. The structure of the product has not been rigorously established, but the structure of the oxygenated product (37) is consistent with elemental analyses and spectral data thus far obtained (Fig. 6).

³M. D. Coburn, J. Heterocyclic Chem. 5, 83 (1968).



REPRODUCED FROM
BEST AVAILABLE COPY

Figure 1. Oxidations of 3-amino-4-(4-chlorophenyl)furan and its sulfilimine and phosphine imine.




Figure 2. Oxidation of aminofurazans and their sulfilimines and phosphine imines with dimethyldioxirane.




Figure 3. Oxidation of 2-aminopyridine and its sulfilimine and phosphine imine with dimethyldioxirane.




Figure 4. Oxidation of aminopyrimidines and their sulfilimines and phosphine imines with dimethyldioxirane.




Figure 5. Oxidation of aminopyrazine and its sulfilimine and phosphine imine with dimethyldioxirane.



Figure 6. Oxidation of 3,6-diamino-1,2,4,5-tetrazine and its disulfilimino derivative.